

| | |
|---------------|--|
| <i>School</i> | <i>Candidate's Name (PLEASE PRINT)</i> |
|---------------|--|



WINCHESTER
COLLEGE

Election

Tuesday 7 May, 2019

Science

PHYSICS

THEORY SECTION

Recommended time: 20 minutes

Write all your answers in the spaces on this question paper

1 To prepare for extra-vehicular activity outside the International Space Station, astronauts train underwater in a pool on Earth called the Neutral Buoyancy Laboratory. An astronaut, in her space-suit, has an average density of 915 kg/m^3 , and a mass of 120 kg (including the suit).


(a) Show that the volume of the suit with the astronaut inside is 0.131 m^3 , stating any formula you use.

.....
..... [1]

(b) In order to simulate some of the effects of 'weightlessness', the astronaut must be made *neutrally buoyant*. This means that her weight must be exactly balanced by the upthrust provided by the water when she is totally immersed. Archimedes' Principle tells us that the *upthrust is equal to the weight of the water displaced by a body*. The pool water has a density of 1.00 g/cm^3 . How much mass must be added to the suit in order to make the astronaut neutrally buoyant? (Assume that the volume of the suit remains constant.)

.....
.....
.....
..... [3]

(c) After having the mass calculated in (b) added to the bottom of her boots, the astronaut is lowered so that she is totally submerged in the water with her body angled at 45° to the vertical. She is then released and finds that she rotates so that her feet point downwards. Use the diagram below to help you explain carefully why this happens, and what the support team would need to do in order to make sure that she remained stationary at 45° to the vertical when released.



.....
.....
.....
.....

(d) The experience of working in the Neutral Buoyancy Laboratory is not identical to that of working in space outside the International Space Station. Suggest one difference that the astronaut might notice.

.....

.....

.....

.....


[1]

2 This question is about the generation of electrical power.

(a) In a nuclear fission power plant, energy is released by splitting up nuclei of uranium-235 atoms. A neutron collides and combines with the uranium-235 nucleus. The nucleus becomes unstable and splits into two large parts, plus some neutrons: this process is called **fission**. Fission releases energy, which is initially carried as the kinetic energy of these two fragments, which is converted to heat energy. This is converted to electrical energy by the turbine and generator.



- (i) An example of a nuclear reaction that can occur in the fission of uranium is shown below:



The masses of the nuclei in this reaction are given below, in atomic mass units, which are represented by the letter u .

$$u = 1.66 \times 10^{-27} \text{ kg.}$$

| | Mass in atomic mass units (u) |
|--------------------------|-----------------------------------|
| ${}_0^1\text{n}$ | 1.008 |
| ${}_{92}^{235}\text{U}$ | 235.048 |
| ${}_{52}^{137}\text{Te}$ | 136.918 |
| ${}_{40}^{97}\text{Zr}$ | 96.906 |

Unlike chemical reactions, the mass can change during nuclear reactions. Show that the change in mass in this reaction is $0.216 u$.

.....

[2]

- (ii) If the mass decreases during a nuclear reaction, the mass deficit is converted to the kinetic energy of the products, using Einstein's famous equation

$$E = mc^2$$

Here $c = 3.00 \times 10^8 \text{ m/s}$ (the speed of light in a vacuum). To use this equation, mass m must be in kilograms, and energy E in joules (J).

1 Calculate the energy released when 6.0×10^{23} uranium-235 nuclei undergo fission by the reaction in (i). This is approximately 235 g of uranium-235.

.....
.....
.....

[3]

2 How long will this number of atoms of uranium-235 keep a power plant with an electrical energy output of 2 GW (2×10^9 joules per second) running? Assume that the power plant converts 33% of the energy released by fission into electrical energy (an efficiency of 33%).

.....
.....
.....

[3]

(iii) When a fission occurs, as well as producing the two large fragments, neutrons are released. These neutrons can go on to cause further fissions. Explain carefully what happens to the reaction if, on average:

1 Fewer than one of these neutrons caused a further fission.

.....
.....
.....
.....

2 Exactly one of these neutrons caused a further fission.

.....
.....
.....
.....

3 More than one of these neutrons caused a further fission.

.....
.....
.....
.....

(b) Coal has a chemical energy density of 39.3 kilojoules per gram (1 kJ = 1000 J). We wish to produce 2.0×10^9 joules of electrical energy per second from our power plant, which is 40% efficient at converting chemical energy in coal to electrical energy. How much coal must be supplied to the power plant per hour?

.....
.....
.....
.....

[4]

End of this paper